

GEL MAX POWER/ GEL NANO

DENTAL WARRANTY DISCLAIMER



CERTIFICATE OF LIMITED WARRANTY AND DISCLAIMER

WARRANTY: Shock Doctor, Inc. ("Shock Doctor") warrants the within mouthguard unit to be free from defects in material and workmanship which could cause damage or injury to the sound, natural teeth of the purchaser herein when properly used in a supervised athletic contest or coach-supervised training session. This warranty shall extend for a period of one (1) year from the date of purchase.

Shock Doctor's obligations under this warranty shall be limited to the lesser of:

- i. The actual dental expense incurred by the user; or
- ii. \$313 per injured or replaced tooth, up to a maximum liability of \$10,000 per incident.

Shock Doctor's obligations hereunder shall be reduced by the amount of any insurance collected or payable for the damage or injury claimed. This certificate does not provide payment or indemnity for orthodontic treatments or preparation of a tooth to be used as an anchor for a bridge

Shock Doctor shall have no responsibility or liability hereunder unless:

- i. The mouthguard was fitted by claimant in accordance with the printed instructions accompanying this warranty; and
- ii. A complete written notice of the injury or damage is received by Shock Doctor within thirty (30) days after the occurrence of the injury or damage; and
- iii. The notice of injury is substantiated by a statement of a licensed dentist providing necessary dental treatment within thirty (30) days after the date of injury or damage and a statement of a participating game official, if a supervised contest, or a supervising coach or school superintendent, if a practice session, that the mouthguard unit was in proper use at the time of injury; and
- iv. The within mouthguard unit is returned to Shock Doctor with the written notice of injury. The mouthguard unit shall become the property of Shock Doctor upon receipt of return.

This warranty is made in lieu of all other warranties, express or implied, including but not limited to the implied warranty for a particular purpose and any implied warranty arising out of a course of a dealing, custom, usage or trade. Except as is otherwise expressly provided herein, Shock Doctor disclaims all liability for loss, injury or damage based upon negligent design or defects in materials and workmanship for the within mouthguard.

Keep this dental warranty, disclaimer and store receipt in your records. In case of injury, mail mouthguard, store receipt, the notice of injury and accompanying statements to Shock Doctor, Inc.

WARNING!

**NOT FOR USE WITH BRACES
DO NOT CHEW YOUR MOUTHGUARD**

In order for a mouthguard to protect, fit, and last it must be worn properly on your teeth and gums as indicated in the fitting instructions. If any mouthguard is chewed or not worn properly, its protection, fit and durability will be greatly diminished.

TO FIT THIS MOUTHGUARD YOU WILL NEED:

- > Mouthguard
- > Cold tap water
- > Saucepan
- > Mirror
- > Boiling water
- > A clock or watch with second hand
- > Slotted spoon

FITTING INSTRUCTIONS:

1. Fill saucepan with 3 to 4 inches of water. Bring water to a boil.
2. Plug the helmet tether strap into front of mouthguard.
3. Remove water from heat source. Let water stand for 30 seconds, then place mouthguard in water for 60 seconds. Do not exceed 60 seconds.
4. Carefully remove mouthguard from hot water with a slotted spoon. Do not remove tether.
5. Cool mouthguard under tap water for 1-2 seconds only to bring surface temperature to a comfortable level.
6. Lick your lips before placing mouthguard in your mouth.
7. Watching in mirror, line up mouthguard carefully with the centerline of upper teeth. Fit mouthguard onto and around upper teeth and press firmly into molars first then front teeth. Do not remove tether.
8. Bring lower jaw forward and up into base of mouthguard with teeth in alignment with lower channel.
9. Close mouth while biting down hard on mouthguard. Suck in strongly and use fingers to press edges of mouthguard onto teeth and roof of mouth through lips and cheeks for 20 seconds.
10. Remove mouthguard and cool under tap water for 30 seconds. Replace into mouth and test for a good, firm fit.
11. Repeat steps 1 through 10 if fit is not accurate.
12. Tether may now be removed.